

RESEARCH ARTICLE

Open access

New geographic records of Harpacticoida (Crustacea: Copepoda) from the Persian Gulf and the Gulf of Oman, Iran

Nazari, F.^{1,2}, Mirshamsi, O.^{1,4*}, Sari, A.³ and Aliabadian, M.^{1,4}

¹Department of Biology, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran

²Department of Biology, Faculty of Science, University of Jiroft, Jiroft, Iran

³School of Biology and Centre of Excellence in Phylogeny of Living Organisms, College of Science, University of Tehran, Tehran, Iran search

⁴Research Department of Zoological Innovations (RDZI), Institute of Applied Zoology, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran

(Received: 4 April 2020; Accepted: 5 July 2020)

Abstract

The meiobenthic harpacticoids of the Persian Gulf and the Gulf of Oman has been almost unknown. During October 2014 to September 2016, interstitial and phytal harpacticoids were collected from tide pools in 30 different localities along the Iranian coast of the Persian Gulf and the Gulf of Oman. The new information on the distribution of 26 species and 12 families is provided, here. The most species-rich family was Laophontidae with seven species, followed by Miracidae with four species. In this paper, all the species were considered new records for the area. Furthermore, the first checklist of meiobenthic Harpacticoids is provided.

Key words: *Harpacticoida*, *Meiobenthics*, *New records*, *Persian Gulf*, *Gulf of Oman*, *Iran*.

INTRODUCTION

Developing information on the biodiversity of copepods in the Persian Gulf and the Gulf of Oman will help to explore changes in the distribution of marine biota of the area. However, there are few published works and taxonomic data that specifically devoted to this taxon in Iranian waters. Of the few studies on Copepods, mostly restricted on zooplankton and meiofauna communities, and also recording parasite copepods (Farhadian & Pouladi, 2014; Taheri-Dezfouli *et al.*, 2016; Hedayati *et al.*, 2017; Sepahvand *et al.*, 2016; Sepahvand *et al.*, 2019). The first study concerned with the marine zooplanktons of the Persian Gulf and the Gulf of Oman, which included marine harpacticoids as well, was performed by Al-Yamani (2011). In this work, four harpacticoid species were reported from the marine zooplankton community of Kuwait waters. Later, the marine harpacticoids were recorded from Iranian waters by Peygan *et al.* (2014). Harpacticoids are as yet unknown from meiofauna and have rarely been reported from planktonic habitat in general. In this study, a preliminary investigation along the Iranian coasts of the Persian Gulf and the Gulf of Oman was performed for documentation of the diversity of meiobenthic harpacticoids. The first contribution concerned with the taxonomy and diversity of meiobenthic copepods of Iran was performed by Nazari *et al.* (2018a, 2018b), three new species and one new record belong to the order Canuella Khodami *et al.* 2017 were identified and described from littoral zone of the Persian Gulf and the

*Corresponding Author: mirshamsi@um.ac.ir



Gulf of Oman. Here, in this contribution, 26 new records of harpacticoids along the Iranian southern coastline are recorded, including the point locality for each species.

MATERIAL AND METHODS

Meiobenthic harpacticoids, reported in this study, were collected from October 2014 to September 2017, from both sediments and seaweeds (Table 1; Fig. 1). Samples were carried out from the littoral zone during low tide using a 60 cc syringe. Sediments were sieved by using sieve with 200 μm and 38 μm mesh size. Macroalga was collected by plastic bag to investigate harpacticoids colonizers. After sorting from sediments and alga, collected specimens were preserved in 96% alcohol for taxonomic investigations. Identifications were aided using a Leica microscope equipped with Differential Interference Contrast (DIC) at 1000 \times magnification and identification keys (Huys *et al.*, 1996; Boxshall & Halsey, 2004; Wells, 2007; Lang, 1948). Moreover, one male and one female of each species were chosen for CLSM photos. Specimens were stained in a mixture of 1:1 Acid Fuchsin and Congo Red overnight (Michels & Büntzow, 2010). Photos were taken by Confocal Laser Scanning Microscope Leica TCS SP5 equipped with a Leica DM500 B. Abbreviations used in the text: EXP, exopod; ENP, endopod; P1-P4, first to fourth swimming legs.




FIGURE 1. Map of sampling sites of the species records in the Persian Gulf and Gulf of Oman.

TABLE 1. Coordinates of the sampling sites and date of sampling.

Map Code	Locality	Longitude	Latitude	Date
1	Mahshahr	30°28'45.17"N	49°11'2.05"E	May 2016
2	Hendijan	30° 6'31.81"N	49°46'20.48"E	May 2016
3	Dylam	30° 2'52.51"N	50° 8'43.97"E	May 2016
4	Bushehr	28°56'44.25"N	50°48'36.86"E	May 2016
5	Halileh	28°50'4.18"N	50°52'32.47"E	May 2016
6	Nayband Bay	27°24'6.66"N	52°37'27.22"E	April 2016
7	North Haleh	27°24'6.12"N	52°38'41.16"E	April 2016
8	Parsian	27° 7'9.05"N	53° 1'15.41"E	April 2016
9	Moqam	27° 40' 01"N	52 ° 19' 54"E	April 2016
10	Bostaneh	26°30'17.16"N	54°38'55.75"E	April 2016
11	Bandar Lengeh	26°34'31.99"N	54°54'52.56"E	April 2016
12	Berke Soflin	26°47'11.79"N	55°19'18.20"E	April 2016
13	Bandar Abbas	27°10'59.01"N	56°19'10.19"E	September 2016
				December 2014
14	Kolahi	27° 2'54.88"N	56°51'0.21"E	October 2015
15	Koohestak	26°48'12.10"N	57° 1'22.33"E	October 2015
				March 2016
16	Sirik	26°31'18.94"N	57° 4'49.91"E	October 2015
17	Old Jask	25°43'48.16"N	57°45'48.99"E	March 2016
18	Jask	25°38'28.01"N	57°46'40.29"E	October 2015
19	Gvanak	25°32'6.35"N	58°52'11.69"E	October 2015
20	Tis	25°21'52.25"N	60°36'29.90"E	October 2015
21	Ramin	25°16'19.94"N	60°44'7.72"E	October 2015
22	Gowatr	25° 9'55.92"N	61°30'6.59"E	October 2015
23	Larak Island	26°53'7.57"N	56°22'57.47"E	April 2015
24	Hangam Island	26°40'12.22"N	55°52'30.33"E	April 2016
25	Hangam Island	26°36'49.11"N	55°52'14.63"E	April 2016
26	Hormoz Island	27° 3'45.66"N	56°30'5.12"E	February 2015
27	Hormoz Island	27° 5'24.97"N	56°28'11.54"E	February 2015
28	Qeshm Island	26°54'47.96"N	56°10'36.29"E	December 2015
29	Qeshm Island	26°41'14.07"N	55°55'55.45"E	December 2015
30	Qeshm Island	26°58'30.42"N	56°14'57.51"E	October 2014

RESULTS

In total, 2748 adult harpacticoids from 26 localities were examined in this study and 26 species belonging to 12 families were identified. The studied taxa are listed here:

Order HARPACTICOIDA Sars, 1903

Family Ameiridae Monard, 1927

Ameira cf. *parvula* (Claus, 1866) (Figs. 2 A-D)

Type. *Canthocamptus parvula* C. Claus 1866, *Schr Ges. Beford. Ges. Naturw. Marburg*. Supplement 9: 1-34.

Distribution of the species. Cosmopolitan. Reported from Germany (Claus, 1866; Pesta, 1932; Kunz, 1935; Lang, 1948); Isle of Man (Moore, 1976); Carolina Island (Vervoort, 1964); North Sea Islands (Mielke, 1975); Andaman and Nicobar Islands (Wells & Rao, 1987); Bulgaria (Apostolov &

Marinov, 1988); Sea of Marmara (Noodt, 1955a; Karaytuğ & Sak, 2006); Mediterranean Sea (Karaytuğ & Sak, 2006; Alper *et al.*, 2010); Korea (Chang, 2007).

Distribution in Iran. Bushehr, Nayband Gulf, North Haleh, Parsian, Bostaneh, Bandar Lengeh, Berke Soflin, Koohestak, Old Jask, Jask, Tis, Larak, Hengam, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Family Dactylopusiidae Lang, 1936

***Dactylopusia tisboides* Claus, 1863 (Figs. 2 E-H)**

Type. *Dactylopus tisboides* C. Claus 1863. *Die freilebenden Copepoden*, pp. 127, t. 16, fig. 24-28. Leipzig.

Distribution of the species. Cosmopolitan species. Mediterranean Sea (Claus, 1863); Nicobar Islands (Sewell, 1940); Sea of Marmara (Noodt, 1955a); Caroline Islands (Vervoort, 1964); Norway (Sars, 1903); British Isles (Brady, 1910); Port Taufiq, Kabret (Gurney, 1927a); Banyuls-Sur-Mer, France (Monard, 1928); Bermuda (Willey, 1935); Australia (Nicholls, 1942); Argentina (Pallares, 1968); Red Sea (Nicholls, 1944); Bulgaria (Apostolov, 1973); Mediterranean Sea (Pulat *et al.*, 2009; Alper *et al.*, 2010).

Distribution in Iran. Kangan, North Haleh, Parsian, Moqam, Bostaneh, Larak, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Diarthrodes cystoecus* Fahrenbach, 1954 (Figs. 2 I-L)**

Type. *Diarthrodes cystoecus* W. H. Fahrenbach, 1954, *J. Wash. Acad. Sci.*, 44: 326.

Distribution of the species. North America (Fahrenbach, 1954, 1962); Tierra del Fuego (Pallares, 1977); Maldive Islands (Sewell, 1940); Madras (Krishnaswamy, 1957); Andaman and Nicobar Islands (Wells & Rao, 1987).

Distribution in Iran. Bushehr, Nayband Gulf, Parsian, Bostaneh, Bandar Lengeh, Berke Soflin, Larak, Hengam, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Paradactylopodia brevicornis* (Claus, 1866) (Fig. 3 A-D)**

Type. *Dactylopus brevicornis* C. Claus, 1866. *Schr. Ges. Beford. Ges. Naturw. Marburg*, suppl. 1:29.

Distribution of the species. Germany (Claus, 1866); Bermuda (Willey, 1935); Nicobar Islands (Sewell, 1940); Sea of Marmara (Noodt, 1955a); Canary Islands (Noodt, 1955b); Karelian coast of White Sea (Chislenko, 1967); Argentina (Pallares, 1975); Andaman and Nicobar Islands (Wells & Rao, 1987); Florida (Walters, 1991); Mediterranean Sea (Pulat *et al.*, 2009).

Distribution in Iran. Bostaneh, Bandar Lengeh, Berke Soflin, Bandar Abbas, Koohestak, Old Jask, Jask, Tis, Larak, Hengam, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Family Harpacticidae Dana, 1846

***Harpacticus ponticus* Marcus, 1967 (Figs. 3 E-H)**

Type. *Harpacticus ponticus* A. Marcus, 1967. *Trav. Mus. Hist. nat. Gr. Antipa*, 7: 29-38, figs. 1-4.

Distribution of the species. Black Sea (Marcus 1967).

Distribution in Iran. Mahshahr, Hendijan, Bushehr, Nayband Gulf, North Haleh, Parsian, Moqam, Berke Soflin, Bandar Abbas, Kolahi, Sirik, Old Jask, Jask, Larak, Hormoz, Hengam, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).




FIGURE 2. Ameiridae: *A. cf. parvula*: A female dorsal, B female ventral, C male dorsal, D male ventral, Dactylopusiidae: *D. tisboides*: E female dorsal, F female ventral, G male dorsal, H male ventral, Dactylopusiidae: *D. cystoecus*: I female dorsal, J female ventral, K male dorsal, L male ventral. Scale: 75 μ m.

Family Laophontidae Scott T., 1904

***Laophonte confusa* Decho & Fleeger, 1986 (Figs. 3 I-L)**

Type. *Laophonte confusa* A. W. Decho & J. W. Fleeger, 1986. *Am. Microsc. Soc.* 105(1): 31-37.

Distribution of the species. Florida Keys (Decho & Fleeger 1986); Gulf of Mexico (Suárez-Morales *et al.* 2009).

Distribution in Iran. Nayband Gulf, North Haleh, Parsian, Moqam, Berke Soflin, Bandar Abbas, Old Jask, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).



FIGURE 3. Dactylopusiidae: *P. brevicornis*: A female dorsal, B female ventral, C male dorsal, D male ventral. Scale: 75 μm .; Harpacticoidae: *H. ponticus*: E female dorsal, F female ventral, G male dorsal, H male ventral. Scale: 100 μm ; Laophontidae: *L. confosa*: I female dorsal, J female ventral, K male dorsal, L male ventral, Scale: 75.

***Laophonte cornuta* Philippi, 1840 (Figs. 4 A-D)**

Type. *Laophonte cornuta* A. Philippi, 1840. *Arch. Naturgesch.* 5: 195.

Distribution of the species. Cosmopolitan. Recorded from Maldiva Archipelago (Sewell, 1933); Red Sea (Nicholls, 1944; Noodt, 1964); Dillon Beach of California (Lang, 1965); Japan (Ito, 1968); Andaman and Nicobar Islands (Wells & Rao, 1987); Belgium (Muller, 2004); China (Liu & Liu, 2008); Gulf of Mexico (Suárez-Morales *et al.*, 2009); New Zealand (Webber *et al.*, 2010); Mediterranean Sea (Pulat *et al.*, 2009).

Distribution in Iran. Bushehr, Nayband Gulf, North Haleh, Parsian, Bostaneh, Berke Soflin, Tis, Gowatr, Larak, Hengam, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Lipomelum heteromelum* Fiers, 1986 (Fig. 4 E-H)**

Type. *Lipomelum heteromelum* F. Fiers, 1986, *Bijd. Dierk.* 56(1).132-164.

Distribution of the species. West Indian Islands, Venezuela, Bahamas and Netherlands Antilles (Fiers 1986).

Distribution in Iran. North Haleh. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Paralaophonte brevirostris* (Claus, 1863) (Figs. 4 I-L)**

Type. *Cleta brevirostris* C. Claus, 1863. *Die Freilebenden Copepodn mit Besonderer Berücksichtigung der Fauna Deutschlands, der Nordsee und des Mittelmeers*: 124.

Distribution of the species. North Sea (Claus, 1863); Chesapeake Bay (Yeatman, 1970); England (Hamond, 1972); Norway (Sars, 1908); Nicobar Islands (Sewell, 1940); Andaman and Nicobar Islands (Wells & Rao, 1987); Mediterranean (Karaytuğ & Sak 2006; Pulat *et al.*, 2009), Black Sea (Kaymak *et al.*, 2012).

Distribution in Iran. Bushehr, Halileh, Kangan, Nayband Gulf, North Haleh, Parsian, Moqam, Bostaneh, Bandar Lengeh, Berke Soflin, Bandar Abbas, Kolahi, Koohestak, Old Jask, Jask, Gvanak, Tis, Gowatr, Larak, Hormoz, Hengam, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Paralaophonte echinata* Fiers, 1986 (Figs. 5 A-D)**

Type. *Paralaophonte echinata* F. Fiers, 1986, *Bijd. Dierk.* 56(1).132-164.

Distribution of the species. West Indian Islands, Venezuela, Bahamas and Netherlands Antilles (Fiers 1986).

Distribution in Iran. North Haleh, Parsian, Berke Soflin, Bandar Abbas, Old Jask, Ramin. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Quinquelaophonte Koreana* Lee, 2003 (Figs. 5 E-H)**

Type. *Quinquelaophonte Koreana* W. Lee, 2003. *Zool. Science.* 20: 657-668.

Distribution of the species. Korea (Lee, 2003)

Distribution in Iran. Ramin, Gowatr. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Quinquelaophonte cf. quinquespinosa* (Sewell, 1924) (Figs. 5 I-L)**

Type. *Laophonte quinquespinosa* R. B. S. Sewell, 1924. *Men. Indian Mus.*, 5: 832.

Distribution of the species. Chilka Lake (Sewell, 1924); Suez canal (Gurney, 1927b); Bermuda (Willey, 1932); Angola (Candeias, 1959); Marseille (Hamond, 1973a); Aldabra (Wells & Meckenzi, 1973); Sirbonian Lagoon (Por, 1973); West Indian Islands, Venezuela, Bahamas and Netherlands Antilles (Fiers,



FIGURE 4. Laophontidae: *L. cornuta*: A female dorsal, B female ventral, C male dorsal, D male ventral; Scale: 100µm. *L. heteromelum*.: E female dorsal, F female ventral, G male dorsal, H male ventral; *P. brevirostris*: I female dorsal, J female ventral, K male dorsal, L male ventral. Scale: 75 µm.




FIGURE 5. Laophontidae: *P. echinata*: A female dorsal, B female ventral, C male dorsal, D male ventral. Scale: 75 μm . *Q. koreana*: E female dorsal, F female ventral, G male dorsal, H male ventral; *Q. cf. quinquespinosa*: I female dorsal, J female ventral, K male dorsal, L male ventral. Scale: 100 μm .

1986); Andaman and Nicobar Islands (Wells & Rao, 1987); Sulawesi (Mielke, 1997); Mexico (Gomez & Morales-Serna, 2013).

Distribution in Iran. Mahshahr, Hendijan, Kolahi. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Family Louriniidae Monard, 1927***Lourinia armata* (Claus, 1866) (Figs. 6 A-D)**

Type. *Jurinia armata* C. Claus, 1866. *Schr. Ges. Beford ges. Naturw. Marburg*, Supplement. 1:30.

Distribution of the species. France, Nice (Claus, 1866); Andaman and Nicobar Islands (Wells & Rao, 1987); Korea (Yoo & Lee, 1993); Mediterranean (Alper *et al.*, 2010).

Distribution in Iran. North Haleh, Nayband Gulf, Parsian, Moqam, Bandar Lengeh, Berke Soflin, Jask, Larak, Hengam, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Family Miraciidae Dana, 1846***Amphiascopsis cf. southgeorgiensis* Lang, 1936 (Figs. 6 E-H)**

Type. *Amphiascopsis southgeorgiensis* K. Lang, 1936. *Further. Zool. Results. Swed. Antarctic. Exped.* 3(3): 1-68.

Distribution of the species. South Georgia Island (Lang, 1936); New Zealand (Hicks, 1971).

Distribution in Iran. North Haleh, Nayband Gulf, Parsian. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Subfamily Diosaccinae Sars M., 1906***Metamphiascopsis hirsutus bermudae* Willey, 1930 (Figs. 6 I-L)**

Type. *Amphiascus hirsutus bermudae* Willey, 1930. *Rep. Govt. Ceylon Pearl Cyster Fish, Gulf Mannar.* 1: 269.

Distribution of the species. Ceylone (Thompson & A. Scott, 1903); Banyuls (Monard, 1928); Bermuda (Willey 1932, 1935); Jamaica (Yeatman, 1976); Nicobar Islands (Sewell, 1940); Caroline Islands (Vervoort, 1964); Andaman and Nicobar Islands (Wells & Rao, 1987); Mediterranean (Sönmez *et al.*, 2014).

Distribution in Iran. Bushehr, North Haleh, Nayband Gulf, Parsian, Bandar Abbas, Gowatr, Larak, Hengam, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Paramphiascella sirbonica* Por, 1973 (Figs. 7 A-D)**

Type. *Paramphiascella sirbonica* F. D. Por, 1973. *Cahiers de Biologie Marine* 14(1): 89-107

Distribution of the species. Sirbonian Lagoon (Por 1973); Gulf of Mexico (Suárez-Morales *et al.* 2009).

Distribution in Iran. Hendijan, Bushehr, Halileh, North Haleh, Nayband Gulf, Parsian, Berke Soflin, Bandar Abbas, Old Jask, Larak, Hengam, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Robertsonia propinqua* Scott T., 1894 (Figs. 7 E-H)**

Type. *Robertsonia propinqua* T. Scott, 1894. *Trans. Linn. Sco. Lond. Sec. 2, 6* (Zool.): 99.

Distribution of the species. Gulf of Guinea (T. Scott, 1894); Chilka Lake (Sewell, 1924); Angola (Candeias, 1959); Argentina (Pallares, 1970); Australia (Hamond, 1973b); Adriatic Sea (Marinov & Apostolov, 1981); Andaman and Nicobar Islands (Wells & Rao, 1987).

Distribution in Iran. Bushehr, North Haleh, Nayband Gulf, Parsian, Bostaneh, Berke Soflin, Bandar Abbas, Old Jask, Kolahi, Koohestak, Sirik, Gowatr, Larak, Hengam, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).



FIGURE 6. Louriniidae: *L. armata*: A female dorsal, B female ventral, C male dorsal, D male ventral; Miraciidae: *A. cf. southgeorgiensis*: E female dorsal, F female ventral, G male dorsal, H male ventral; *M. hirsutus bermuda*: I female dorsal, J female ventral, K male dorsal, L male ventral. Scale: 100 μ m.




FIGURE 7. Miraciidae: *P. sirbonica*: A female dorsal, B female ventral, C male dorsal, D male ventral. Scale: 75 μ m. *R. propinqua*: E female dorsal, F female ventral, G male dorsal, H male ventral. Scale: 100 μ m. Parastenheliidae: *P. hornelli*: I female dorsal, J female ventral, K male dorsal, L male ventral; *P. cf. spinosa*: M female dorsal, N female ventral, O male dorsal, P male ventral. Scale: 75 μ m.

Family Parastenheliidae Lang, 1936

***Parastenhelia hornelli* Thompson I. C. & Scott A., 1903 (Figs. 7 I-L)**

Type. *Parastenhelia hornelli* I. C. Thompson & A. Scott, 1903, *Rep. Govt. Ceylon Pearl Oyster Fish. Gulf Manaar*, 1:263.

Distribution of the species. Cosmopolitan. Recorded from: Sri Lanka (Thompson & A. Scott, 1903); Carolina Islands (Vervoort, 1964); New Zealand (Wells *et al.*, 1982); Mozambique (Wells, 1967); Caribbean Sea, Barbados (Coull, 1970); Virgin Isles (Coull, 1971, Hartzband & Hummon, 1974); Black Sea (Apostolov, 1973); Sea of Marmara (Noodt, 1955a); Andaman and Nicobar Islands (Wells & Rao, 1987).

Distribution in Iran. Bushehr, North Haleh, Nayband Gulf, Parsian, Bostaneh, Hormoz. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Parastenhelia cf. spinosa* (Fischer, 1860) (Figs. 7 M-P)**

Type. *Harpacticus spinosus* S. Fischer, 1860, *Asb. Bayer. Akad. Wiss., Abt. 3*, 8: 665.

Distribution of the species. Cosmopolitan. Recorded from: Plymouth area of England (Fraser, 1936); Maldive Archipelago (Sewell, 1940); Helgoland (Klie, 1941); Canary Island (Noodt, 1955b);

Caroline Islands (Vervoort, 1964); Bulgaria (Apostolov, 1968); Andaman and Nicobar Islands (Wells & Rao, 1987); Mediterranean Sea (Apostolov, 1973).

Distribution in Iran. Bushehr, North Haleh, Nayband Gulf, Parsian, Moqm, Bandar Lengeh, Berke Soflin, Jask, Ramin, Larak, Hengam, Hormiz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Family Peltidiidae Claus, 1860

***Peltidium* cf. *perplexum* Thompson I. C & Scott A., 1903 (Figs. 8 A-D)**

Type. *Peltidium perplexum* I. C. Thompson & A. Scott, 1903, *Rep. Govt. Ceylon Pearl Oyster Fish. Gulf Manaar*, 1:263.

Distribution of the species. Ceylone (Thompson & A. Scott 1903).

Distribution in Iran. Larak, Hengam, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Family Pseudotachidiidae Lang, 1936

***Idomene* cf. *laticaudatus* Thompson I. C & Scott A., 1903 (Figs. 8 E-H)**

Type. *Dactylopusia laticauda* Thompson & A. Scott, 1903. *Rep. Govt. Ceylon Pearl Oyster Fish. Gulf Manaar*, 1: 263.

Distribution of the species. Ceylone (Thompson & A. Scott 1903); Maldiv Archipelago (Sewell, 1940); Inhaca Island, Mozambique (Wells, 1967).

Distribution in Iran. Tis. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Family Tegastidae Sars G. O., 1904

***Syngastes parilis* Bartsch, 1994 (Figs. 9 A-C)**

Type. *Syngastes parilis* I. Bartsch, 1994. *Spixiana, Munich*, 17, 2:161-173.

Distribution of the species. Australia (Bartsch 1994).

Distribution in Iran. Bushehr, Nayband Gulf, Parsian, Bostaneh, Larak, Hengam, Hormoz, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

***Tegastes* cf. *clausi* Sars, 1904 (Fig. 9D)**

Type. *Tegastes longimanus* Claus, 1863. *sensu* G. O. Sars, 1904. *An Account of the Crustacea of Norway Bergen Muse*, 5:57-80.

Distribution of the species. North Sea (Claus, 1863); West coast of Norway (Sars, 1904); Banyuls-Sur-Mer, France (Monard, 1928); Helgoland (Klie, 1949).

Distribution in Iran. Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Family Thalestridae Sars G. O., 1905

***Eudactylopus* cf. *latipes* (Scott T., 1893) (Figs. 10 A-D)**

Type. *Dactylopus latipes* T. Scott, 1893. *Transact. Linn. Soc. London. 2nd Series: Zool.* 6(1): 1-161.

Distribution of the species. Gulf of Guinea (Scott T., 1894); Nicobar Islands, the Ceylon Pearl Banks (Thompson & A. Scott, 1903); Malay Archipelago (A. Scott, 1909); Nicobar Island and Maldiv Archipelago (Sewell, 1940); Marmara Sea (Noodt, 1955a); Shingu (Tanaka & Hue, 1967).

Distribution in Iran. Nayband Gulf, North Haleh, Parsian, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

Family Tisbidae Stebbing, 1910

***Tisbe alaskensis* Chullasorn, Dahms, Lee, Ki, Schizas, Kangtia, Park & Lee, 2011 (Figs. 10 E-H)**

Type. *Tisbe alaskensis* S. Chullasorn, H. Dahms, K. Lee, J. Ki, N. Schizas, P. Kangtia, H. G. Park & J. Lee, 2011. *Zool. Studi.* 50, 1: 103-117.

Distribution of the species. Juneau, Alaska (Chullasorn *et al.*, 2011).

Distribution in Iran. Bushehr, Parsian, Tis, Ramin, Qeshm. New to the Persian Gulf and the Gulf of Oman (Fig. 1).




FIGURE 8. Peltidiidae: *P. cf. perplexum*: A female dorsal, B female ventral, C male dorsal, D male ventral. Scale: 100µm. Pseudotachidiidae: *I. cf. laticaudatus*: E female dorsal, F female ventral, G male dorsal, H male ventral. Scale: 50µm.



FIGURE 9. Tegastidae: *S. parilis*: A female lateral, B male lateral, C male grasping female; D *T. cf. clausi*: female lateral. Scale: 50 μ m.

***Tisbintra jonesi* Ummerkutty, 1960 (Figs. 10 I-L)**

Type. *Tisbintra jonesi* N. P Ummerkutty, 1960. *Mar. Biol. Ass. India*, 2, 2:149-164

Distribution of the species. Gulf of Mannar (Ummerkutty, 1960)

Distribution in Iran. Bandar Abbas, Kolahi, Old Jask. New to the Persian Gulf and the Gulf of Oman (Fig. 1).

DISCUSSION

In this study, we present a total of 26 species belonging to 12 harpacticoid families. All taxa are reported for the first time from the Persian Gulf and the Gulf of Oman. These were compared carefully with original species descriptions. However, the identification of some species has to be defined with reservation (names indicated with cf.).

Ameira parvula is represented as the species complex based on variation in the morphological characters in different populations (See Wells, 2007). Thus, revision of the species based on morphological and molecular studies is necessary.

Dactylopusia tisboides was mentioned as a cosmopolitan species (Vervoort, 1964). It has been reported from all oceans in the world, but only a few published works have been shown the distribution and description of the species. Wells (2007) mentioned the species as either variable cosmopolitan or complex. Therefore, the decision must be taken with caution.

Quinquelaophonte quinquespinosa is a cosmopolitan species. There is interpopulation variability in the number of antennary exopod seta, number of setae on P4 exp-3 and enp-2, P1 exp-2: exp-1 length ratio, furcal rami length: width and the number of female P5 armatures (Wells & McKenzie, 1973; Lee, 2003). Therefore, this was considered as a species complex by Lee (2003). The material was found in Iran were congruent well with the descriptions presented by Lee (2003) and Gomez & Morales-Serna (2013), especially female which was described by the later authors (they found only the female from Mexico). Using *COI* barcoding is the only way to clear species delimitation of this species.

Moreover, the decision about *A. cf. southgeorgiensis*, *P. cf. spinosa*, *P. cf. perplexum*, *I. cf. laticaudatus*, *T. cf. clause* and *E. cf. latipes* is uncertain. Unfortunately, insufficient information on these taxa hampered definite identification of the species. Furthermore, in some cases, species delimitation can be possible only by using molecular data. Therefore, poor genetic data makes the positions of these taxa unclear.

The result of this survey is only a small part of copepods diversity in the Persian Gulf and the Gulf of Oman. High species diversity and richness in copepod fauna indicated in this study is in its early stage and further thorough taxonomic surveys are needed to shed more light to our current knowledge on this taxon in the region.

ACKNOWLEDGMENTS

We would like to thank Professor Pedro Martinez Arbizu the headmaster of the German Center for Marine Biodiversity Research (DZMB) and Dr. Kai Horst George (DZMB) for their valuable comments and verifications of species identifications. Also, authors have special thanks to Dr. Gritta Veit-Kohler for providing access to DZMB laboratory facilities. This study was supported by a grant from the Office of Research Affairs, Ferdowsi University of Mashhad, Iran to H.B. (Project 3/32569).

LITERATURE CITED

Alper, A., Karaytuğm, S., Sak, S., 2010. Interstitial and phytal Harpacticoida (Crustacea: Copepoda) inhabiting the mediolittoral zone of the Datça-Bozburun Peninsulas (Muğla, Turkey). *SDU Journal of Science* 5, 16–28.

Al-Yamani, F.Y., Skryabin, V., Gubanov, A., Khvorov, S., Prusova, I., 2011. Marine zooplankton practical guide for the northwestern of Arabian Gulf. *Kuwait Institute for Scientific Research* 2, 1–210.

Apostolov, A., Marinov, T., 1988. Copepoda Harpacticoida (Morski harpaktikoidi). *Fauna of Bulgaria* 18, 1–383.




FIGURE 10. Thalestridae: *E. cf. latipes*: A female dorsal, B female ventral, C male dorsal, D male ventral. Scale: 100µm. Tisbodae: *T. alaskensis*: E female dorsal, F female ventral, G male dorsal, H male ventral. Scale: 75µm. *T. jonsi*: I female dorsal, J female ventral, K male dorsal, L male ventral. Scale: 100 µm.

Apostolov, A., 1968. Neue und bemerkenswerte harpacticoide Ruderfußkrebse (Crustacea Copepoda) aus dem Küstengrundwasser Bulgariens. Zoologischer Anzeiger 180, 395-402.

Apostolov, A., 1973. Apport vers l'étude d'Harpacticoïdes pontiques habitant les algues marines. Zoologischer Anzeiger 191, 263-28.

Bartsch, I., 1994. Three new species of *Syngastes* from south-western Australia (Tegastidae, Harpacticoida, Copepoda). Spixiana, Munich 17, 161-173.

Boxshall, G.A., Halsey, S.H., 2004. An introduction to copepod diversity. The Ray Society, London, pp. 1-966.

Brady, G.S., 1910. Die marinen Copepoden der Deutschen Südpolar Expedition 1901-1903. I, Ueber die Copepoden der Stämme Harpacticoida, Cyclopoida, Notodelphyoida und Caligoida. Deutsch. Südpolar Expedition, 1901-1903, 11, Zoologie 3, 497-594.

Candeias, A., 1959. Contribution to the knowledge of the harpacticoids (Crustacea, Copepoda) from the littoral of Angola. Publicações Culturais da Companhia de Diamantes de Angola (Diamang) 45, 79-104.

Chang, C.Y., 2007. Two harpacticoid species of genera *Nitokra* and *Ameira* (Harpacticoida: Ameiridae) from brackish water in Korea. Integrative Biosciences 11, 247-253.

Chislenko, L.L., 1967. Copepoda Harpacticoida of the Karelian coast of the White Sea. Garpaktitsidy (Copepoda Harpacticoida) Karelskogo poberezh'ya Belogo morya. In: Hidrobiol. Issled. na Karel'skom poberezh'e Belogo morya. Issledovaniya Fauny Morei 7, 48-196.

Chullasorn, S., Dahms, H., Lee, K., Ki, J., Schizas, N., Kangtia, P., Park, H.G., Lee, J., 2011. Description of *Tisbe alaskensis* sp. nov. (Crustacea: Copepoda) Combining Structural and Molecular Traits. Zoological Studies 50, 103-117.

Claus, C., 1863. Die frei lebenden Copepoden mit besonderer Berücksichtigung der Fauna Deutschlands, der Nordsee und des Mittelmeeres. Verlag von Wilhelm Engelmann, Leipzig, 1-230.

Claus, C., 1866. Die Copepoden-Fauna von Nizza. Ein Beitrag zur Charakteristik der Formen und deren Abänderungen im Sinne Darwins. Schriften der Gesellschaft zur Beförderung der Gesamten Naturwissenschaften zu Marburg, Supplement 1, 1-34.

Coull, B.C., 1970. Harpacticoid copepods from Barbados and Jamaica, W.I., with descriptions of two new species. Caribbean Journal of Science 10, 129-135.

Coull, B.C., 1971. Meiobenthic Harpacticoida (Crustacea, Copepoda) from St. Thomas, U.S. Virgin Islands. Transactions of the American Microscopical Society 90, 207-218.

Decho, A.W., Fleeger, J.W., 1986. A new meiobenthic species of *Laophonte* (Copepoda: Harpacticoida) from the Florida Keys. Transactions of the American Microscopical Society 105, 31-37.

Fahrenbach, W.H., 1954. A new species of the genus *Diarthrodes* (Crustacea, Copepoda) parasitic in a red alga. Journal of the Washington Academy of Sciences 44, 326-329.

Fahrenbach, W.H., 1962. The biology of a harpacticoid copepod. La Cellule 62, 303-376.

Farhadian, O., Pouladi, M., 2014. Seasonal changes in the abundance and biomass of zooplankton from shallow mudflat river-estuarine system in the Persian Gulf. Aqua Science and Technology 18, 19-29.

Fiers, F., 1986. Harpacticoid copepods from the West Indian Islands: Laophontidae (Copepoda, Harpacticoda). Amsterdam Expedition to the West Indian Islands, Report 48. Bijdr Dierkd 56, 132-164.

Fischer, S., 1860. Beiträge zur Kenntniss der Entomostraceen. Abhandlungen der Bayerischen Akademie der Wissenschaften. Mathematisch - Naturwissenschaftliche Abteilung, 8, Abt, 3, 645-682.

Fraser, J.H., 1936. The distribution of rock pool Copepoda according to tidal level. *Journal of Animal Ecology* 5, 23-28.

Gomez, S., Morales-Serna, F.N., 2013. On a small collection of Laophontidae T. Scott (Copepoda: Harpacticoida) from Mexic. II. New records of *Quinquelaophonte* Wells, Hiks and Coull and description of *Onychoquinpes permixtionis* gen. nov. et sp. nov. *Journal of Natural History* 47, 5-12, DOI:10.1080/00222933.2012.75765.

Gurney, R., 1927a. Zoological results of the Cambridge expedition to the Suez Canal, 1924 23. Report on the Crustacea-Copepoda (littoral and semi-parasitic). *Transactions of the Zoological Society of London*, 22, 451-577.

Gurney, R., 1927b. Report on the Crustacea. Copepoda of brine-pools at Kabret. Zoological results of the Cambridge Expedition to the Suez Canal, 1924. *Transactions of the Zoological Society of London* 22, 173-177.

Hamond, R., 1972. Some marine and brackish-water copepods from Wells-next-the-Sea, Norfolk, England. *Transactions of the Norfolk and Norwich Naturalists Society* 22, 237-243.

Hamond, R., 1973a. The harpacticoid copepods (Crustacea) of the saline lakes in southeast Australia, with special reference to the Laophontidae. *Records of the Australian Museum* 28, 393-420.

Hamond, R., 1973b. The Australian species of *Robertsonia* (Crustacea, Harpacticoida), with a revised key to the genus. *Records of the Australian Museum* 28, 421-435.

Hartzband, D.J., Hummon, W.D., 1974. Sub-community structure in subtidal meiobenthic Harpacticoida. *Oecologia, Berlin* 14, 37-51.

Hedayati, A., Pouladi, M., Vazirizadeh, A., Qadermarzi, A., Mehdipour, N., 2017. Seasonal variations in abundance and diversity of copepods in Mod River estuary, Bushehr, Persian Gulf. *Biodiversitas* 18, 447-452.

Hicks, G.R.F., 1971. Some littoral harpacticoid copepods, including five new species, from Wellington, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 5, 86-119.

Huys, R., Gee, J.M., Moore, C.G., Hamond, R., 1996. Marine and brackish water harpacticoids, Part 1. *Synopses of the British Fauna*, No. 51. Field Studies Council, Shrewsbury, pp. 1-352.

Karaytuğ, S., Sak, S., 2006. A contribution to the marine Harpacticoid (Crustacea, Copepoda) fauna of Turkey. *Ege Journal of Fisheries and Aquatic Sciences* 23, 403-405.

Kaymak, N.B., Karaytuğ, S., Sak, S., 2012. Laophontidae fauna (Crustacea: Copepoda: Harpacticoida) of the Turkish Black Sea coast. *Journal of Anatolian Natural Sciences* 3, 23-36.

- Khodami, S., McArthur, J.V., Blanco-Bercial, L., Martinez Arbizu, P., 2017. Molecular phylogeny and revision of Copepod orders (Crustacea: Copepoda). Scientific Reports 7(1), 9164. DOI:10.1038/s41598-017-06656.
- Klie, W., 1941. Marine Harpacticoiden von Island. Kieler Meeresforschungen 5, 1-44.
- Klie, W., 1949. Harpacticoida (Cop.) aus dem Bereich von Helgoland und der Kieler Bucht. 1. Kieler Meeresforschungen 6, 90-128.
- Krishnaswamy, S., 1957. Studies on the Copepoda of Madras. Thesis, University of Madras, 1-168.
- Kunz, H., 1935. Zur Ökologie der Copepoden Schleswigholsteins und der Kieler Bucht. Schr naturw Ver Schlesw-Holst 21, 84- 132.
- Lang, K., 1936. *Copepoda Harpacticoida*. In: Bock, S. (ed.). Further Zoological Results of the Swedish Antarctic Expedition, 1901-1903. Stockholm 3, 1-68.
- Lang, K., 1948. Monographie der Harpacticiden. Nordiska-Bokhandeln, Stockholm, 2 Vols, pp. 1-1682.
- Lang, K., 1965. Copepoda Harpacticoida from the Californian Pacific coast. Kungliga Svenska Vetensk-Akademiens Handlingar, Fjarde Serien. Almqvist & Wiksell, Stockholm 10, 1-560.
- Lee, W., 2003. A marine harpacticoid, *Quinquelaophonte koreana* sp. nov. from a sandy beach in Korea (Crustacea: Copepoda). Zoological Sciences 20, 657-668.
- Liu, R., Liu, J.Y., 2008. Checklist of marine biota of China seas. China Science Press. 1267 pp.
- Marcus, A., 1967. Harpacticoids from the rocky facies in the Black Sea: *Harpacticus ponticus* n. sp., *Tisbe dilatata* Klie, *Ectinosoma melaniceps* Boeck, *Halectinosoma herdelongata* n.n. (syn. *Ectinosoma intermedium* Marcus). Travaux du Muséum National d'Histoire Naturelle Grigore Antipa 7, 29-38.
- Marinov, T., Apostolov, A., 1981. Contribution à l'étude des Copépodes Harpacticoides de la mer Adriatique (côte Yougoslave). 2. Sur le méiobenthos du cap Piran. Acta Zoologica Bulgarica 18, 23-30.
- Michels, J., Büntzow, M., 2010. Assesment of Congo red as a flurescence maker for the exoskeleton of small crustaceans and the cuticle of polychates. Journal of Microscopy 238, 95-101.
- Mielke, W., 1975. Systematik der Copepoda eines Sandstrandes der Nordseeinsel Sylt. Mikrofauna Meeresbodens 52, 1-134.
- Mielke, W., 1997. On a small collection of Laophontidae (Copepoda) from Sulawesi, Indonesia. Microfauna Marina 11, 223-250.
- Monard, A., 1928. Les harpacticoids marins de Banylus. Archives de zoologie expérimentale et générale 67, 259-443.

Moore, C.G., 1976. The harpacticoid families Thalestridae and Ameiridae (Crustacea, Copepoda) from the Isle of Man. *Journal of Natural History* 10, 29-56.

Muller, Y., 2004. Faune et flore du littoral du Nord, du Pas-de-Calais et de la Belgique: inventaire. [Coastal fauna and flora of the Nord, Pas-de-Calais and Belgium: inventory]. Commission Régionale de Biologie Région Nord Pas-de-Calais: France, 307 pp.

Nazari, F., Mirshamsi, O., Sari, A., Aliabadian, M., 2018a. A first report of *Canuellina insignis* Gurney, 1927 (Canuellida: Copepoda) from the Persian Gulf and Gulf of Oman. *Iranian Journal of Animal Biosystematics* 14, 131-136.

Nazari, F., Mirshamsi, O., Sari, A., Aliabadian, M., & Martinez Arbizu, P. 2018b. Three new Canuellidae Copepoda: Canuelloida) from Iran. *Zootaxa* 4446, 401– 441.

Nicholls, A.G., 1942. Marine Copepoda from Western Australia. I. Littoral Harpacticoids from Rottnest Island. *Journal of the Royal Society of Western Australia* 27, 135-141.

Nicholls, A.G. 1944. Littoral Copepoda from the Red Sea. *The Annals and magazine of natural history*, 11, 487-503.

Noodt, W., 1955a. Marmara Denizi Harpacticoid'leri (Crust. Cop.). Marine Harpacticoiden (Crust. Cop.) aus dem Marmare Meer. *Istanbul Universitesi Fen Fakultesi Mecmuasi* 20, 49-94.

Noodt, W., 1955b. Copepoda Harpacticoidea von Teneriffa (Kanarische Inseln). *Zoologischer Anzeiger* 154, 200-222.

Noodt, W., 1964. Copepoda Harpacticoidea aus dem Litoral des Roten Meeres. *Kiel Meeresforschungen* 20, 128-154.

Pallares, R.E., 1977. Copépodos harpacticoides marinos de Tierra del Fuego (Argentina). Isla de los Estados II. El género *Diarthrodes* Thomson, 1882. *Contribuciones Científicas CIBI- MA, Contribuciones Científicas del Centro de Investigaciones de Biología Marina (CIBIMA)*, Buenos Aires 141, 1-13.

Pallares, R.E., 1968. Copépodos marinos de la Ría Deseado (Santa Cruz, Argentina). *Contribución sistemática-ecológica. II. Physis*, Buenos Aires 27, 245-262

Pallares, R.E., 1970. Copépodos Marinos de la Ría Deseado (Santa Cruz, Argentina). *Contribución Sistemático-Ecológica. III. Physis*, Buenos Aires 30, 255-282.

Pallares R.E., 1975. Copepods harpacticoides marinos de Tierra del Fuego (Argentina). 1. Isla de los Estados. *Contribuciones Científicas del Centro de Investigaciones de Biología Marina (CIBIMA)*, Buenos Aires 141, 1-13.

Pesta, O., 1932. Krebstiere oder Crustacea. 1. Ruderfüsser oder Copepoda. 3. Unterordnung: Harpacticoida (1 and 2). In: Dahl, F. (ed.). *Die Tierwelt Deutschlands und der Angrenzenden Meeresteile nach Ihren Merkmalen und nach Ihrer Lebensweise*. Jena 24, 1-164.

- Peyghan, S., Savari, A., Doustshenas, B., Sakhaee, N., Dehghan, S., 2011. New record of *Acartia* (*Acartiella*) *faoensis* Khalaf, 1991 (Copepoda: Calanoida: Acartidae) from Iranian waters of NW Persian Gulf. Iranian Journal of Animal Biosystematics 7, 177-179.
- Philippi, A., 1940. Zoologische Bemerkungen (Fortsetzung). IV. Kurze Charakteristik mehrerer neuer Genera aus der Familie der Copepoden. Archiv der Natur 6, 181-195.
- Por, F.D., 1973. The benthic Copepoda of the Sirbonian Lagoon (Sabkhat el Bardawil). Cahiers de Biologie Marine 14, 89-107.
- Pulat, İ., Özel, İ., Aker, V., 2009. Gümüldür sahili (Ege Denizi) mediolittoral kayalık biyotoplarından tespit edilen Thalestridae ve Laophontidae (Copepoda, Harpacticoida) türleri. Ege Journal of Fisheries and Aquatic Sciences 26, 55-58.
- Sars, G.O., 1903. An Account of the Crustacea of Norway, with short descriptions and figures of all the species. Copepoda, Calanoida, with supplement. Bergen Museum, Alb. Cammermeyer's Forlag, Christiana 4, 1-171.
- Sars, G.O., 1904. Copepoda Harpacticoida. Parts V & VI. Harpacticidae (concluded), Peltidiidae, Tegastidae, Porcellidiidae, Idyidae (part). An Account of the Crustacea of Norway, with short descriptions and figures of all the species. Bergen Museum 5, 57-80.
- Sars, G.O., 1908. Copepoda Harpacticoida. Parts XXI & XXII. Laophontidae (continued). An Account of the Crustacea of Norway, with short descriptions and figures of all the species. Bergen Museum, Bergen 5, 241-256.
- Scott, A., 1909. The Copepoda of the Siboga Expedition. Part I. Free-swimming, littoral and semi-parasitic Copepoda. Siboga Expeditie, Monograph, Leiden 29, 1-323.
- Scott, T., 1894. I. Report on Entomostraca from the Gulf of Guinea, collected by John Rattray, B.Sc. Transactions of the Linnean Society of London. 2nd Series: Zoology 6, 1-161.
- Sepahvand, V., Kihara, T.C., Boxshall, G.A., 2019. A new species of *Clausidium* Kossmann, 1874 (Copepoda: Cyclopoida) associated with ghost shrimps from the Persian Gulf, including female-male interlocking mechanisms and remarks on host specificity. Systematic Parasitology 96, 171-189. DOI 10.1007/s11230-019-09839-x.
- Sepahvand, V., Rastegar-Pouyani, N., Kihara, T.C., Momtazi, F., 2017. A new species of *Clausidium* Kossmann, 1874 (Crustacea, Copepoda, Cyclopoida, Clausidiidae) associated with ghost shrimps from Iran. Journal of the Brazilian Crustacean Society, DOI 10.1590/2358-2936e2017018.
- Sewell, R.B.S., 1924. Fauna of the Chilka Lake. Crustacea Copepoda. Memoirs of the Indian Museum, Calcutta 5, 771-851.
- Sewell, R.B.S., 1933. Notes on a small collection of marine Copepoda from the Malay States. The Raffles Bulletin of Zoology 8, 25-31.
- Sewell, R.B.S., 1940. Copepoda, Harpacticoida. Scientific Reports of John Murray Expeditions 7, 117-382.

Sönmez, S., Sak, S., Karaytuğ, S., 2014. Marine interstitial and phytal Miraciidae Dana, 1846 (Crustacea: Copepoda: Harpacticoida) inhabiting along the mediolittoral zone of Turkish coasts. *Journal of Anatolian Natural Sciences* 5, 52–96.

Suárez-Morales, E., Fleeger J.W., Montagna, P.A., 2009. Free-Living Copepoda (Crustacea) of the Gulf of Mexico, Pp. 841–869 in Felder, D.L. and D.K. Camp (eds.), *Gulf of Mexico–Origins, Waters, and Biota. Biodiversity*. Texas A&M University Press, College Station, Texas.

Taheri-Dezfouli, T., Nabavi, S.M.B., Rajabzadeh-Ghatromi, E., Sajjadi, N., 2016. Study on the Meiofauna Community Structure in Sajafi Shore as the Bio-Indicator of Environmental Population. *Scientific Research Publishing* 6, 632–644.

Tanaka, O., Hue, J.S., 1967. Preliminary report on the copepods found in the tide pool along the north-west coast of Kyushu. *Proceedings of the Symposium on Crustacea, Ernakulam*, 12–15 January 1965. *Symposium Series, Journal of the Marine Biological Association of India* 2, 57–73.

Thompson, I.C., Scott, A., 1903. Report on the Copepoda collected by Professor Herdman, at Ceylon, in 1902. In: Herdman, W.A. (ed.). *Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar*. 1, suppl. 7, 227–307.

Ummerkutty, A.N.P., 1960. Studies on Indian copepods. 2. An account of the morphology and life history of a harpacticoid copepod, *Tisbintra jonesi* sp. nov. from the Gulf of Mannar. *Journal of the Marine Biological Association of India* 2, 149–164.

Vervoort, W., 1964. Free-Living Copepoda from Ifaluk Atoll in the Caroline Islands with Notes on related species. *Bulletin of the United States National Museum* 236, 1–431.

Walters, K., 1991. Influences of abundance, behavior, species composition, and ontogenetic stage on active emergence of meiobenthic copepods in subtropical habitats. *Marine Biology* 108, 207–215.

Webber, W.R., Fenwick, G.D., Bradford-Grieve, J.M., Eagar, S.G., Buckeridge, J.S., Poore, G.C.B., Dawson, E.W., Watling, L., Jones, J.B., Wells, J.B.J., Bruce, N.L., Ahyong, S.T., Larsen, K., Chapman, M.A., Olesen, J., Ho, J., Green, J.D., Shiel, R.J., Rocha, C.E.F., Lörz, A., Bird, G.J., Charleston, W.A., 2010. Phylum Arthropoda Subphylum Crustacea: shrimps, crabs, lobsters, barnacles, slaters, and kin. In: Gordon, D.P. (Ed.) (2010). *New Zealand inventory of biodiversity: 2. Kingdom Animalia: Chaetognatha, Ecdysozoa, Ichnofossils* pp. 98–232.

Wells, J.B.J., 2007. An annotated checklist and keys to the species of Copepoda Harpacticoida (Crustacea). *Zootaxa* 1568, 1–872.

Wells, J.B.J., McKenzie, K.G., 1973. Report on a small collection of benthic copepods from marine and brackish waters of Aldabra, Indian Ocean. *Crustaceana* 25, 133–146.

Wells, J.B.J., Rao, G.C., 1987. Littoral harpacticoida (Crustacea: Copepoda) from Andaman and Nicobar Islands. *Zoological Survey of India* 16, 1–385.

Wells, J.B.J., 1967. The littoral Copepoda (Crustacea) of Inhaca Island, Mozambique. *Transactions of the Royal Society of Edinburgh* 67, 189–358.

Wells, J.B.J., Hicks, G.R.F., Coull, B.C., 1982. Common harpacticoid copepods from New Zealand harbours and estuaries. *New Zealand Journal of Zoology* 9, 151-184.

Willey, A., 1930. Harpacticoid copepods from Bermuda. Part I. *The Annals and magazine of natural history, Series 10*, 15, 50-100.

Willey, A., 1932. Copepod phenology – observations based on new material from Canada and Bermuda. *Atti del Congresso Internazionale Zoologica*, 1930. *Archives Zoology Italiana* 16, 601-617.

Willey, A., 1935. Harpacticoid Copepoda from Bermuda. Part II. *The Annals and magazine of natural history, Series 10*, 50-100.

Yeatman, H.C., 1970. Copepods from Chesapeake Bay sponges including *Asterocheres jeanyeatmanae* n. sp. *Transactions of the American Microscopical Society* 89, 27-38.

Yeatman, H.C., 1976. Marine littoral copepods from Jamaica. *Crustaceana (Leiden)* 30, 201-219.

Yoo, K., Lee, W., 1993. A marine harpacticoid, *Lourinia armata* (Claus, 1866) New to Korea (Crustacea: Copepoda). *Korean Journal of Systematic Zoology* 9, 115-121.